

Final Version [1.2]

**Richmond upon Thames
Suicide and Self-Harm
Prevention Strategy**

2019 - 2022

Richmond upon Thames Council

Richmond CCG 2020



Executive Summary	2
Vision.....	2
Aim.....	2
Objectives.....	2
1. Where are we now?	3
1.1 Richmond upon Thames Profile.....	3
1.2 What is the Scale of the Problem?	4
1.3 Self-Harm.....	5
1.4 Suicide Audit 2014	5
1.5 Suicide Prevention Framework	8
2 Where do we want to be?	9
2.1 Vision.....	9
2.2 Aim.....	9
2.3 Objectives.....	9
2.4 Outcomes	9
2.5 Enablers.....	10
2.6 Guiding Principles	10
2.7 Approach	10
2.8 Risk Factors/ Groups.....	11
2.9 Community Action Plan	12
3 How will we get there?.....	14
3.1 Strategic Objectives and Action Plan	14
4 How Will We Know When We've Got There? - Monitoring the Strategy.....	17
4.1 Governance.....	17
4.2 Interdependent Action Plans	17
4.3 Baseline Metrics 18/19 and beyond	17

Executive Summary

Vision

Our vision is that Richmond will be a home to communities that are described as happy, thriving and resilient.

We believe that with the right support at the right time, people can recover from psychological distress and mental disorder.

We believe suicide is preventable and aim to ensure individuals;

- value their own life and the lives of others
- should never feel that suicide is the only option
- are supported in times of need, by safe, integrated, and compassionate services

Aim

The overall aim of this strategy is to reduce the rate of suicide, suicidal behaviour and self-harm through the following objectives:

Objectives

This strategy aims to:

- 1. Improve understanding of local need** – Suicide audit, pathway mapping and needs assessment
- 2. Challenge the stigma and discrimination associated with mental disorder by:**
 - Increasing awareness of mental health and mental disorder
 - Adopt more sensitive media reporting in relation to mental disorder, self-harm and suicide
- 3. Improve access to information and Postvention Support** - for those concerned or affected by suicide
- 4. Prevent self-harm amongst young people** – Support the development of the CAMHS Local Transformation Plan including prevention in schools, effective care pathways and A&E liaison services
- 5. Improve access to services** - Increase the numbers of people from high risk groups accessing appropriate services
- 6. Improve crisis responses and pathways** – Improvements in crisis care pathways and closer integration of mental health and substance misuse services, safer discharge, improved crisis planning.

Governance

This Strategy has been agreed by the Crisis Care Concordat and approved by the Health and Wellbeing Board. The Crisis Care Concordat will be responsible for overseeing the progress and delivery of the action plan and will report back to the Health and Well-being Board on an annual basis.

1. Where are we now?

1.1 Richmond upon Thames Profile

1.1.1 The table below can be used to see how Richmond compares to London on several measures and risk factors associated with suicide. (Public Health England, 2020)

Compared with benchmark: ● Better ● Similar ● Worse ○ Not compared ● Lower ● Similar ● Higher ○ Not compared

Quintiles: Best ● ● ● ● ● Worst Low ● ● ● ● ● High ○ Not applicable

Recent trends: - Could not be calculated ➔ No significant change ↑ Increasing / Getting worse ↑ Increasing / Getting better ↓ Decreasing / Getting worse ↓ Decreasing / Getting better ↑ Increasing ↓ Decreasing


[Export table as image](#) [Export table as CSV file](#)

Indicator	Period	Richmond		Region England			London region		
		Recent Trend	Count	Value	Value	Value	Worst/Lowest	Range	Best/Highest
Depression: Recorded prevalence (aged 18+)	2018/19	↑	13,378	7.2%	7.6%*	10.7%*	5.7%		10.3%
Mental Health: QOF prevalence (all ages)	2018/19	➔	2,066	0.87%	1.12%*	0.96%*	0.73%		1.51%
Estimated prevalence of opiate and/or crack cocaine use	2016/17	-	623	4.9	9.3	8.9	14.4		4.9
Long-term health problem or disability: % of population	2011	-	21,447	11.5%	14.2%	17.6%	17.3%		11.2%
Self-reported wellbeing - people with a low satisfaction score	2018/19	-	-	*	4.1%	4.3%	-	Insufficient number of values for a spine chart	-
Self-reported wellbeing - people with a low worthwhile score	2018/19	-	-	*	3.4%	3.6%	-	Insufficient number of values for a spine chart	-
Self-reported wellbeing - people with a low happiness score	2018/19	-	-	*	7.3%	7.8%	-	Insufficient number of values for a spine chart	-
Self-reported wellbeing - people with a high anxiety score	2018/19	-	-	20.8%	21.0%	19.7%	29.0%		15.3%
Prisoner population: count	Sep 2018	-	-	-	7677*	78533*	-	Insufficient number of values for a spine chart	-
Children in the youth justice system (10-18 yrs)	2017/18	↓	88	2.4*	5.9	4.5	12.6		2.4
Children leaving care: rate per 10,000 children aged under 18	2017/18	➔	64	14.2	27.3	25.2	14.2		160.6
Marital breakup: % of adults	2011	-	15,271	10.2%	10.6%	11.6%	12.8%		7.7%
Social Isolation: percentage of adult social care users who have as much social contact as they would like (18+ yrs)	2018/19	-	830	47.9%	41.3%	45.9%	33.9%		53.5%
Social Isolation: percentage of adult carers who have as much social contact as they would like (18+ yrs)	2018/19	-	25	21.0%	33.2%	32.5%	16.7%		45.7%
Older people living alone: % of households occupied by a single person aged 65 or over	2011	-	9,434	11.8%	9.6%	12.4%	6.0%		13.9%
People living alone: % of all usual residents in households occupied by a single person	2011	-	26,008	14.1%	12.8%	12.8%	8.0%		23.4%
Unemployment (model-based)	2018	-	4,100	3.8%	5.0%	4.1%	6.3%		3.6%
Statutory homelessness - Eligible homeless people not in priority need	2017/18	-	-	*	1.0	0.8	2.8		0.1
Statutory homelessness - households in temporary accommodation	2017/18	➔	282	3.3	14.9	3.4	40.1		1.9
Domestic abuse-related incidents and crimes	2018/19	-	-	32.9*	32.9	27.4	32.9		32.9
Long term claimants of Jobseeker's Allowance	2018	➔	335	2.8	4.2	3.8	8.2		1.0
Children in care	2019	➔	115	25	50	65	86		25
Admission episodes for alcohol-related conditions (Broad) (Persons)	2018/19	↑	3,467	1,989	2500	2367	3,379		1,854
Admission episodes for alcohol-related conditions (Broad) (Male)	2018/19	↑	2,254	2,868	3523	3246	4,899		2,535
Admission episodes for alcohol-related conditions (Broad) (Female)	2018/19	↑	1,213	1,277	1637	1608	2,182		1,235
Estimated prevalence of common mental disorders: % of population aged 16 & over	2017	-	20,430	13.2*	19.3*	16.9*	24.4		13.2
Estimated prevalence of common mental disorders: % of population aged 65 & over	2017	-	2,514	8.3*	11.3*	10.2*	14.6		8.3

1.2 What is the Scale of the Problem?

- 1.2.1 Every suicide is both an individual tragedy and a terrible loss to society, it affects a number of people directly and often many others indirectly. The impact can be devastating –psychologically, economically, and spiritually – for all those affected. Suicides can be prevented, mainly by having a society that is caring and compassionate, that supports people at times of personal crisis and ensures that marginalised individuals are helped.
- 1.2.2 There are a wide range of local stakeholders that have a role to play in preventing suicide. Many people who die by suicide have not been in touch with mental health services. There are many things we can do in our communities and outside hospital and care settings to help those who think suicide is their only option.
- 1.2.3 Suicide is a major issue for society and a leading cause of years of life lost. Although the number of people who take their own lives in England has followed a downward trend over the last 30 years, the suicide rate in England is currently 10.1 per 100,000 compared to 8.2 per 100,000 in London. The suicide rate in Richmond, 9.2 per 100,000, remains below the England average but is higher than the London average this rate equates to 10–15 suicides per year.

1.2.4 **Figure 2: Number of suicides and death due to injury of undetermined intent in Richmond by gender, all ages (3 year average).**



Source: Public Health England. 2020. Suicide Prevention Profile.

1.3 Self-Harm

- 1.3.1 Self-harm is when somebody intentionally damages or injures their body. It's usually a way of coping with or expressing overwhelming emotional distress. More than half of people who die by suicide have a history of self-harm. Although some people who self-harm are at a high risk of suicide, most people who self-harm do not want to end their life. Self-harm is a risk factor for suicide and provides an important indicator to inform preventative interventions. A recent commentary of suicide prevention research concluded that effective interventions for reducing self-harm in both adolescents and the adult population remains our most effective tool in preventing suicide (Sedgwick and Ougrin, 2019)¹.
- 1.3.2 Since 2011, hospital admissions for self-harm in young people aged 10–24 are consistently higher than the London average. The most recent data (2016/17) identifies that Richmond has the highest rate of self-harm (10–24) in London at 433.6 per 100,000, this equates to 116 individuals per year.
- 1.3.3 In 2014 public health carried out a Self-Harm Audit. During 2012/13–2013/14, self-harm accounted for 0.4% of A&E attendances and 1.1% of hospital admissions. The majority of self-harm admissions were due to self-poisoning (92%) and 6% were recorded as self-harm by a sharp object. The highest numbers and rates of both self-harm related A&E attendances and hospital admissions were in females age 15–24, although males aged 20–24 also experienced relatively high rates of self-harm admissions. A smaller peak was identified in females aged 50–54 for A&E attendances. There is some evidence that the white British ethnic group made up a greater than expected proportion of self-harm attendances and admissions.
- 1.3.4 Home was the location of the incident leading to most self-harm hospital admissions (87%). Relatively high numbers of self-harm A&E attendances and hospital admissions came from North Sheen/South Kew and Whitton areas and relatively low numbers came from the area of St. Margaret's and East Twickenham. Eleven per cent of people attending A&E for deliberate self-harm left without receiving treatment. Comparing the number of admissions from A&E to hospital for self-harm between the two datasets suggests self-harm is particularly under-recorded in A&E.

1.4 Suicide Audit 2014

- 1.4.1. Deaths included in the 2014 audit include those given a verdict of suicide, open and narrative (including some drug related deaths) by the Coroner. Statistics on causes of death produced by the Office for National Statistics (ONS) are based on the information provided at death registration. These statistics are provided to the Department of Health on an annual basis. Open verdicts are generally coded by the ONS as deaths from injury or poisoning of undetermined intent. When national statistics are presented, suicides and deaths of undetermined intent are combined.

¹ Sedgwick, R., Ougrin, D., (2019). No better than chance? Developments in predicting adolescent suicide, a commentary on Mars et al. (2018) and Beckman et al. (2018). *The Journal of Child Psychology and Psychiatry*, Volume 60, Issue 1 January 2019.

This reflects research studies which show that the majority of open verdicts are most likely suicides, although they do not meet the high legal standard of evidence required for a Coroner to record a suicide verdict. Therefore, official suicide rates are measured using this broader definition and are reflected in this suicide prevention strategy.

- 1.4.2. A Suicide audit provides intelligence to inform strategic priorities. Results from the Richmond Suicide audit further confirm this. Audits are time consuming requiring detailed interrogation and close working with the Coroner. We are currently exploring opportunities to develop real-time data collection which will reduce the need for resource intensive retrospective audits in the future. Thrive London are in the process of piloting a real time pan-London database to quickly identify suspected suicides. We propose to utilise this database to improve our surveillance of local suicides. In the interim the 2014 audit combined with national findings will inform the ongoing development of the strategy.
- 1.4.3. According to the Primary Care Mortality Database extract supplied by NHS Digital, between 2006 and 2013, 94 suicides (death due to intentional self-harm or poisoning/injury of undetermined intent) in Richmond residents were registered; an average of 12 per year
- 1.4.4.
- Thirty-seven (39%) of the suicides registered between 2006 and 2013 in Richmond were in females and 57 (61%) were in males. This is more equal than may be expected: nationally, males account for about three quarters of all suicides.
 - Over half the deaths (52%) occurred in the 35–54-year-old age group.
 - One-third of suicides were due to poisoning, 31% were due to hanging, strangulation and suffocation and 36% were due to other methods
 - The most common marital status for those included in the audit was single, at 40% of the total.
 - The single commonest living situation for those included in the audit was living alone (32%).
 - The single most common employment status was employed (30%) followed by retired (22%), unemployed (18%) and not known (12%).
 - A physical health problem was recorded in 51% of cases and there was no significant difference in men (52%) and women (50%).
 - Three-quarters of the cohort were recorded as having any history of mental illness. Women were slightly more likely to have been recorded as having problems with mental health than men (77% versus 72%).
 - In total, 39% of all cases were recorded as having a history of previous self-harm. Men were more likely to have a history of self-harm than women (43% versus 34%)
 - 42% of cases in the audit had a history of problems with alcohol. Men were significantly more likely to have had problems with alcohol than women (56% versus 23%).

- 27% of cases were reported to have had a history of substance misuse. Men had higher rates of substance misuse than women (34% versus 16%).
- 15% of cases were explicitly recorded as having financial difficulties at the time of death.
- Over a fifth (22%) of cases were found to have had relationship difficulties at the time of death.
- 27% of individuals included in this audit had experience of violence, excluding a history of self-harm or violence to oneself. Men were more likely to have been recorded as having experienced violence (30% versus 23% of women).
- Sixteen per cent of individuals included in the audit had a recorded history of criminal offence, incarceration, probation service involvement or were suspects in a criminal investigation. The proportion was much higher in males (25%) than females (5%)

1.5 Suicide Prevention Framework

1.5.1 The framework was developed in 2015 and sets out priorities across six key areas. It takes account of the evidence about what measures are likely to be effective for reducing suicide and self-harm, (including the National Institute for Health and Care Excellence guidance) and key elements have been included in the action plan included in this strategy. The framework was developed based upon discussions at a multi-agency workshop (September 2014) involving those working in mental health, community safety, substance misuse, social care, primary care, and children's services and also a number of service users and representatives.

1.5.2 Six priority areas:

- Reduce the risk of suicide in key high-risk groups
- Tailor approaches to improve mental health among specific groups
- Prevent self-harm among young people
- Reduce access to the means of suicide
- Provide better information and support to those bereaved or affected by suicide
- support research, data collection and monitoring

1.5.3 The Suicide Prevention Framework 2015 provides a solid foundation to inform the next stage of our local strategy. The priorities outlined above are aligned to the national strategy and are consistent with the strategic objectives set out in this strategy. In many ways this strategy provides an evolution of the earlier framework and includes updated recommendations from the national strategy.

2 Where do we want to be?

2.1 Vision

Our vision is that Richmond will be a home to communities that are described as happy, thriving and resilient.

We believe that with the right support at the right time, people can recover from psychological distress and mental disorder.

We believe suicide is preventable and aim to ensure individuals;

- value their own life and the lives of others
- should never feel that suicide is the only option
- are supported in times of need, by safe, integrated, and compassionate services

2.2 Aim

The overall aim of this strategy is to reduce the rate of suicide, suicidal behaviour and self-harm through the following objectives:

2.3 Objectives

This strategy aims to:

1. **Improve understanding of local need** – Suicide audit, pathway mapping and needs assessment
2. **Challenge the stigma and discrimination associated with mental disorder by:**
 - Increasing awareness of mental health and mental disorder
 - Adopt more sensitive media reporting in relation to mental disorder, self-harm and suicide
3. **Improve access to information and ‘Postvention’ support** – for those concerned or affected by suicide
4. **Prevent self-harm amongst young people** – Support the development of the CAMHS Local Transformation Plan including prevention in schools, effective care pathways and A&E liaison services
5. **Improve access to services** – Increase the numbers of people from high risk groups accessing appropriate services
6. **Improve crisis responses and pathways** – Improvements in crisis care pathways and closer integration of mental health and substance misuse services, safer discharge, improved crisis planning.

2.4 Outcomes

- Reduction in suicides – maintain a suicide rate below the London average
- Positive experience of care
- Increased use of safety planning in key services
- Recovery and staying well

2.5 Enablers

- A multi-agency partnership approach to deliver a defined action plan (Crisis Care Concordat)
- Feedback and consultation with stakeholders
- Focus on primary and secondary prevention
- Clinical Commissioning support for Strategic Actions
- Seamless and integrated primary / secondary care pathways
- Focus on local intelligence gathering and surveillance

2.6 Guiding Principles

All activities undertaken as part of this strategy should be guided by the following principles:

Principle	Description
Equity and Equality	Provision of services should be proportional to need to avoid increasing health inequalities and targeted to areas that need them the most.
Accessibility	Services should be accessible to all, with factors including, geography, opening hours and access for people with disabilities considered.
Early Intervention and Prevention	People will receive information, opportunities and support to help them take care of their own health and wellbeing, prevent deterioration and lead independent lives.
Integration	Service provision and care pathways should be integrated, with all relevant providers working together. This will maximise the benefits of delivery
Effectiveness	Services should be evidence-based and provide value for money
Quality	Services and activities commissioned will be of high quality, with quality and patient/ resident playing a key role in the assessment of what makes a 'good' service. The delivering of quality services will be a high-profile factor in the commissioning process.
Sustainability	Services should be developed and delivered with consideration of social, economic and environmental sustainability.
Safeguarding	People health and social care services will be safeguarded, throughout their experience.
Dignity and Respect	All people who use or encounter services will be treated with dignity and respect, recognising that people are often going through difficult times.

2.7 Approach

The Richmond Crisis Care Concordat (CCC) was formed in 2016 and provides a multi-agency forum to lead on improved crisis care for people experiencing psychological distress. The development and implementation of the strategy and action plan will be the responsibility of this group. Key partners in the delivery of this strategy are; Richmond upon Thames Borough Council, Achieving for Children, Richmond Clinical Commissioning Group, Primary Care, South West London & St. George's Mental Health Trust, East London Foundation Health Trust, Metropolitan Police, service users, families/carers, communities and the voluntary sector.

2.8 Risk Factors/ Groups

Risk Groups	Risk Factors	Additional Risk factors	Mental Health Conditions²
Young and middle-aged men	Life History; Trauma in childhood; history of sexual or physical abuse; history of parental neglect	Being gay, lesbian or transgender, arising from the prejudice these groups often face	Severe depression; People with severe depression are much more likely to attempt suicide than the general population
People in the care of mental health services	Mental health; developing a serious mental health condition	Being in debt	Bipolar disorder; About one in three people with bipolar disorder will attempt suicide at least once. People with bipolar disorder are 20 times more likely to attempt suicide than the general population
History of self-harm and suicide	Drugs or alcohol misuse	Being homeless	Schizophrenia; It is estimated that 1 in 20 people with schizophrenia will take their own life.
People in the criminal justice system	Employment; poor job security, low levels of job satisfaction or being unemployed	Being a war veteran	Borderline personality disorder; It is estimated that just over half of people with borderline personality disorder will make at least one suicide attempt.
Specific occupational groups including doctors, veterinary workers, farmers and agricultural workers	Stressful life event; Bereavement, separation long term / terminal health condition	Being in prison or recently released from prison	Anorexia nervosa; It is estimated that around one in five people with anorexia will make at least one suicide attempt.
	Genetics and Family history	Occupations that provide access to potential ways of dying by suicide e.g. working as a doctor, nurse, pharmacist, farmer or as a member of the armed forces	

² NHS Choices, <http://www.nhs.uk/Conditions/Suicide/Pages/Causes.aspx>

2.9 Confidential Enquiry into Suicide by Children and Young People

2.9.1 The purpose of a confidential enquiry is to detect areas of deficiency in clinical practice and devise recommendations to resolve them. During 2014 and 2015, 922 suicides by people aged under 25 in England and Wales were examined.

The enquiry found:

- The number of suicides at each age rose steadily in the late teens and early 20s.
- Most of those who died were male (76%), and the male to female difference was greater in those over 20.
- Academic pressures and bullying were more common before suicide in under 20s, while workplace, housing and financial problems occurred more often in 20–24 year olds.

2.9.2 The following themes were identified in the national report;

- Bereavement was common in both age groups, 25% of under 20s and 28% of 20–24 year olds, equivalent to around 125 deaths per year.
- Suicide bereavement was more common in the under 20s (11% v 6%)
- University and college students accounted for 21% of under 20s and 14% of 20–24 year olds. Only 12% were reported to be seeing student counselling services.
- Looked After Children accounted for 9% of under 20s who died with high rates of housing problems and suicidal ideation, one third of the cohort were not in recent contact with mental health care.
- LGBT young people accounted for 6% of under 20s and 3% of 20–24 year olds; a quarter of LGBT under 20 had been bullied; most had previously self-harmed.
- Suicide related internet use was implicated in 26% of deaths in under 20s, and 13% of deaths in 20–24 year olds
- Self-harm was reported in 52% of under 20s and 41% of 20–24 year olds who died. Around 60% in both age groups were known to services and around 40% had been in recent contact—in only 26% this was mental health care.
- Interagency collaboration was variable and risk recognition appeared poor.

2.9.3 The following key messages were identified in the report;

- Suicide in young people is rarely caused by one thing; it usually follows a combination of previous vulnerability and recent events.
- The stresses identified that lead to suicide are common in young people; most come through them without serious harm.
- Important themes for suicide prevention are support for or management of family factors childhood abuse, bullying, physical health, social isolation, mental ill-health and alcohol or drug misuse.
- Specific actions are needed for the following groups (1) support for young people who are bereaved, especially by suicide (2) greater priority for mental health in colleges and universities (3) housing and mental health care for looked after children (4) mental health support for LGBT young people.

- Further efforts are needed to remove information on suicide methods from the internet; and to encourage online safety, especially for under 20s.
- Suicide prevention in children and young people is a role shared by front-line agencies; they need to improve access, collaboration and risk management skills.
- Services which respond to self-harm are key to suicide prevention in children and young people and should work with services for alcohol and drug misuse, factors that are linked to subsequent suicide.

2.10 Community Action Plan

2.10.1 During 2018, a task group, led by public health, was set up to formulate a Community Action Plan to mitigate against the risk of Suicide Clusters and contagion. A small multi-agency group created the plan based upon a practice resource developed by Public Health England.

2.10.2 The term “suicide cluster” describes a situation in which more suicides than expected occur in terms of time, local, or both. A suicide cluster usually includes three or more deaths. With suicidal behaviour increasingly spreading via the internet and social media, a greater number of suicides than expected may well occur in a specific time period and be spread out geographically (so called mass clusters).

2.10.3 Suicide clusters understandably cause great concern and may lead to hasty responses. It is important that plans for such occurrences are prepared in advance, to ensure a measured and effective response. With this in mind the Richmond Suicide Community Action Plan provides a systematic process to identify potential clusters and provide a series of swift interventions to mitigate the threat.

2.10.4 The Community Action Plan has two specific facets: active surveillance through a Suicide Surveillance Group and mitigation through a multi-agency Suicide Response Group. The plan has been developed with partners including Public Health, Clinical Commissioning Group, Metropolitan Police, Local Safeguarding Children Board, Adult Social Care, Achieving for Children. The Suicide Surveillance Group will meet regularly throughout the year to monitor the levels of suicide and to identify potential suicide clusters and contagion.

3 How will we get there?

3.1 Strategic Objectives and Action Plan 2019/20

Objective	Action	Measure / Outcome	Owner	Timescale
1. Understand local need	1.1 Set up a system to monitor suicide through a Suicide Surveillance Group	Quarterly meeting of the Suicide Surveillance Group	GM	Sept 19
	1.2 Develop a Suicide Prevention survey for Primary Care (GPs and Practice Nurses) to ascertain levels of need related to self-harm and suicide.	Survey developed, disseminated and recommendations drafted	CCG AD	Mar 20
2. Challenge stigma and Discrimination	2.1 Distribute suicide reporting guidelines to local media groups	Media organisations identified, and guidance distributed with accompanying letter from Director Public Health	GM	Sept 19
	2.2 Mental Health First Aid and Suicide Prevention Training delivered to Frontline staff and Community and Voluntary Sector	Baseline established, and annual summary of courses delivered	GM	Mar 20
3. Improve access to Information and 'Postvention' support	3.1 Support Secondary state maintained and independent schools to access suicide prevention training.	20 Schools accessing Papyrus suicide prevention training for schools	GM	April 19
	3.2 Primary Care fact sheet distributed to primary care settings (GP practices and health centres)	Primary care fact sheet distributed to GP practices and health centres	CCG AD	Mar 20
	3.3 Help is at hand booklets and Z cards distributed to Primary Care settings	All GP and Health Centres provided with Help is at Hand materials	CCG AD	Sept 19
	3.4 Suicide prevention signage at high risk locations including railway stations, bridges and level crossings	Location of additional signage reported to CCC on quarterly basis	HR	Mar 20
4. Prevent self-harm amongst young people	4.1 Promote whole school approach to emotional health and well-being as set out in the CAMHS LTP	Whole school approach audit implemented, and best practice seminars developed and attended by schools	CCG DR	Sept 19

	4.2 Ensure support for young people who self-harm through locally agreed self-harm pathway	Self-harm pathway developed and disseminated to universal services	CCG DR	Mar 20
5. Improve Access to local services	5.1 Improve the Integrated support for individuals experiencing co-occurring mental health and substance misuse support	Clinical Interface meetings to receive recommendations on cases identified through the Drug and Alcohol Related Deaths Panel.	RK	Dec 19
	5.2 Improve access to psychological therapies for key risk groups including men.	Key risk groups identified and KPIs agreed with commissioner and provider	CCG AD	Mar 20
	5.3 Develop and implement Suicide Community Action Plan to mitigate against cluster and contagion.	Community Action Plan agreed by Crisis Care Concordat and first meeting of Suicide Surveillance Group to monitor suicide data	GM	Sept 19
	5.4 Increase access to Mental Health First Aid Training and suicide prevention training for school's staff, first responders and universal services	Baseline established, and annual summary of courses delivered.	GM	Mar 20
6. Improve crisis responses and pathways	6.1 Review crisis pathways across Richmond by mapping access point to treatment and supporting and ensuring co-ordinated and integrated care across primary and secondary mental health services in response to need	CCG to facilitate the crisis care mapping which will form part of the South West London crisis work. Local task and finish group to link into the South West London Crisis work.	CCG AD	Mar 20
	6.2 Ensure safer hospital discharge to include follow up within 72 hours after leaving hospital.	95% of service users leaving hospital will have a face to face review and risk assessment within 72 hrs of discharge.	CHa	Mar 20
	6.3 Review of the SWL pathways for Children and young people requiring crisis.	Healthy London Partnership to undertake a peer review of SWLSTG crisis care pathways.	CCG DR	Dec 19
	6.4 Effective liaison in urgent care centres and A&E.	95% of attendances to urgent care, receiving an assessment from the Core 24 team within 1 hr. KPI monthly monitoring	CCG AD	Dec 19

4 How Will We Know When We've Got There? - Monitoring the Strategy

4.1 Governance

4.1.1 The Suicide Prevention Strategy has been agreed by the Richmond Crisis Care Concordat; The Health and Wellbeing Board and the Adult Social Services, Health and Housing Services Committee.

4.1.2 The multi-agency Richmond Crisis Care Concordat will be responsible for overseeing the progress and delivery of the action plan and will report back to the Health and Well-being Board annually.

4.2 Interdependent Action Plans

- CAMHS Transformation Plan 2018
- LSCB Young Peoples Risky Behaviour Action Plan 2018
- SWL Health and Care Partnership Plan 2019
- South West London and St. George's Mental Health Trust Suicide Prevention Plan

4.3 Baseline Metrics 18/19 and beyond

4.3.1 The following table identifies key indicators

Indicator	Baseline (18/19)	19/20
Suicide Rate	7.9 Per 100,000	9.2 Per 100,000
Emergency Hospital Admission for Intentional Self-Harm	132.2 per 100,000	Awaiting data update
Number of Richmond patients in secondary care dying by suicide	5	Awaiting data update
Adult males accessing psychological therapies	1,313	Awaiting data update
No of MHFA training sessions	18	Awaiting data update

BIBLIOGRAPHY

Cross-Government Suicide Prevention Workplan, published in January 2019 by the Department of Health and Social Care

<https://www.gov.uk/government/publications/suicide-prevention-cross-government-plan>

Future in mind: promoting, protecting and improving our children and young people's mental health and wellbeing, Published by the Department of Health and NHS England in 2012

www.gov.uk/government/uploads/system/uploads/attachment_data/file/414024/Childrens_Mental_Health.pdf

Local Suicide Prevention Planning, A practice resource, Published by Public Health England in 2016

<https://www.gov.uk/government/publications/suicide-prevention-developing-a-local-action-plan>

Preventing Suicide in England: a cross-government outcomes strategy to save lives, Published by the department of Health in 2012

<https://www.gov.uk/government/publications/suicide-prevention-strategy-for-england>

Preventing suicide in England: Third progress report of the cross-government outcomes strategy to save lives, published by the Department of Health and Social Care in 2017

<https://www.gov.uk/government/publications/suicide-prevention-third-annual-report>

Preventing suicide in England: fourth progress report of the cross-government outcomes strategy to save lives, published by the Department of Health and Social Care in 2019

<https://www.gov.uk/government/publications/suicide-prevention-fourth-annual-report>

Suicide prevention: identifying and responding to suicide clusters, published by Public Health England in 2015

<https://www.gov.uk/government/publications/suicide-prevention-identifying-and-responding-to-suicide-clusters>

Crisis Care Concordat Membership

Organisations

Richmond Clinical Commissioning Group
Achieving for Children
East London Foundation Trust
LBRuT (Adult Social Services)
LBRuT (Public Health)
LBRuT (Housing)
LBRuT (Safeguarding Adults Team)
Richmond GPs
Richmond Mind
London Ambulance Service
SW London Health and Care Partnership
Metropolitan Police
British Transport Police
Network Rail
South West Trains
Comfort Care/ Retreat Crisis House
SWLStG Mental Health Trust
SPEAR
Child and Adolescent Mental Health Services
Substance Misuse Services (CGL)